

Systematic review

BJS Open

Open Access

Prevention of incisional hernia after midline laparotomy with prophylactic mesh reinforcement: a meta-analysis and trial sequential analysis

A. P. Jairam¹ , M. López-Cano⁴ , J. M. Garcia-Alamino⁶, J. A. Pereira⁵, L. Timmermans², J. Jeekel³, J. Lange³ and F. Muysoms⁷

¹Department of Surgery, Catharina Hospital Eindhoven, Eindhoven, and ²Department of Surgery, Maasstad Ziekenhuis Rotterdam and ³Department of General Surgery, Erasmus University Medical Centre, Rotterdam, the Netherlands, ⁴Department of General and Digestive Surgery, Hospital Vall d'Hebrón, Universitat Autònoma de Barcelona, and ⁵Department of General and Digestive Surgery, Hospital Del Mar, Barcelona, Spain, ⁶Department of Primary Health Care Sciences, University of Oxford, Oxford, UK, and ⁷Department of Surgery, Maria Middelares Hospital, Ghent, Belgium

Correspondence to: Dr A. P. Jairam, Catharina Hospital Eindhoven, Michelangelolaan 2, 5623 EJ Eindhoven, the Netherlands (e-mail: anjairam@hotmail.com)

Background: Incisional hernia is a frequent complication after abdominal surgery. The aim of this study was to assess the efficacy of prophylactic mesh reinforcement (PMR) after midline laparotomy in reducing the incidence of incisional hernia.

Methods: A meta-analysis was conducted following PRISMA guidelines. The primary outcome was the incidence of incisional hernia after follow-up of at least 12 months. Secondary outcomes were postoperative complications. Only RCTs were included. A random-effects model was used for the meta-analysis, and trial sequential analysis was conducted.

Results: Twelve RCTs were included, comprising 1815 patients. The incidence of incisional hernia was significantly lower after PMR compared with sutured closure (risk ratio (RR) 0.35, 95 per cent c.i. 0.21 to 0.57; $P < 0.001$). Both onlay (RR 0.26, 0.11 to 0.67; $P = 0.005$) and retromuscular (RR 0.28, 0.10 to 0.82; $P = 0.02$) PMR led to a significant reduction in the rate of incisional hernia. The occurrence of seroma was higher in patients who had onlay PMR (RR 2.23, 1.10 to 4.52; $P = 0.03$). PMR did not result in an increased rate of surgical-site infection.

Conclusion: PMR of a midline laparotomy using an onlay or retromuscular technique leads to a significant reduction in the rate of incisional hernia in high-risk patients. Individual risk factors should be taken into account to select patients who will benefit most.

[Correction added on 19 February 2020, after first online publication: J. García Alamino has been amended to J. M. Garcia-Alamino]

Funding information

No funding

Paper accepted 17 December 2019

Published online 14 February 2020 in Wiley Online Library (www.bjsopen.com). DOI: 10.1002/bjs.50261

Introduction

Incisional hernia (IH) is a frequent complication after abdominal surgery, with an incidence ranging from 11 to 20 per cent in general surgical populations^{1–4}. The incidence of IH can increase up to 40 per cent in high-risk groups, such as patients with an abdominal aortic aneurysm (AAA) or morbid obesity^{5–12}. IH may be asymptomatic, but can also lead to serious and potentially fatal complications, such as incarceration and strangulation of bowel. Furthermore, IH has a high impact on patients' quality of

life and body image^{13,14}, and treatment of IH represents a financial burden on the healthcare system¹⁵.

Current treatment of IH is mesh repair, which has led to a lower recurrence rate compared with the primary suture technique¹⁶. However, the recurrence rate is still high, even when mesh is used. A Danish nationwide registry study¹⁷ reported a cumulative recurrence rate of 37 per cent at 3 years after IH repair. Currently, there is no definitive solution for the high recurrence rates and complications related to these recurrences. Prevention is therefore of paramount importance¹⁸. In the past

Fig. 1 PRISMA diagram for the review




Table 1 Risk-of-bias assessment for prevention of incisional hernia with prophylactic mesh reinforcement in midline laparotomy

Reference	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)
Abo-Ryia <i>et al.</i> ⁹	–	–	–	–		+
Ball <i>et al.</i> ¹⁰	–	–	–	–	+	+
Bevis <i>et al.</i> ⁸	+	+	+	+	+	+
Caro-Tarrago <i>et al.</i> ²⁹	+		+	+		+
El-Khadrawy <i>et al.</i> ²⁸	–	–	–	–		+
García-Ureña <i>et al.</i> ³¹	+		+	+	+	+
Gutiérrez de la Peña <i>et al.</i> ⁶	–	–	–	–		+
Jairam <i>et al.</i> ²³	+	+	+	+		+
Muysoms <i>et al.</i> ¹¹	+	+	+	+		+
Pans <i>et al.</i> ⁵	–	–	–	–		+
Sarr <i>et al.</i> ³⁰	+	–	–	–	–	+
Strzelczyk <i>et al.</i> ⁷	+		+	+		+

+, Low risk of bias; –, high risk of bias.

Table 2 Summary of findings for included studies on the prevention of incisional hernia with prophylactic mesh reinforcement in midline laparotomies



Reference	n	Indication for midline laparotomy	Type of mesh	Mesh position	Follow-up (months)	Outcome measurements	Diagnosis of IH (clinical/radiological)
Pans <i>et al.</i> ⁵	288	Morbid obesity	Polyglactin	Intraperitoneal	29.8	IH, postoperative morbidity	Not reported
Gutiérrez de la Peña <i>et al.</i> ⁶	88	Colorectal cancer, gastric cancer, cholelithiasis, diverticulosis, Crohn's disease, pancreatic cystadenoma, gastric ulcer, cancer of small intestine	Polypropylene	Onlay	36	IH, haematoma, seroma, infection, pain	Clinical; if not conclusive, CT
Strzelczyk <i>et al.</i> ⁷	74	Gastric bypass surgery	Polypropylene	Retromuscular	28	IH, wound leak, bleeding, other surgical complication	Clinical plus ultrasound imaging
El-Khadrawy <i>et al.</i> ²⁸	40	High risk	Polypropylene	Preperitoneal	36.7	IH, seroma	Clinical
Bevis <i>et al.</i> ⁸	80	AAA	Polypropylene	Retromuscular	Mesh 30.2 No mesh 19.6	SSI, wound disruption, chronic wound pain, cardiac, pulmonary problems, DVT, ascites IH, wound infection, hernia operation	Clinical; if doubt, ultrasound imaging
Abo-Ryia <i>et al.</i> ⁹	64	Open bariatric surgery	Polypropylene	Preperitoneal	Mesh 48 No mesh 49	Safety and efficacy of preperitoneal prosthetic enforcement, seroma, infection, partial dehiscence	Clinical; ultrasound imaging in suspected cases
Caro-Tarrago <i>et al.</i> ²⁹	160	Colorectal and general surgery	Polypropylene	Onlay	Mesh 14.8 No mesh 12.5	IH, all adverse events, postoperative complications	Clinical and CT
Sarr <i>et al.</i> ³⁰	280	Open RYGB	Biological	Intraperitoneal	24	IH, wound infection, wound dehiscence, wound sinus tract, wound erythema, seroma	Clinical, phone call, primary care physician
Bali <i>et al.</i> ¹⁰	40	Open AAA	Biological	Onlay	36	IH, duration of surgery, postoperative complications, reoperation rate	Clinical and CT
García-Ureña <i>et al.</i> ³¹	107	Colorectal surgery	Polypropylene	Onlay	24	IH, incidence of local complications: SSI, seroma, evisceration, mesh rejection	Clinical and CT
Muysoms <i>et al.</i> ¹¹	114	AAA and ASA grade < IV	Partially absorbable polypropylene	Retromuscular	24	Systemic complications IH	Clinical and, if available, ultrasound imaging or CT
						Postoperative complications SSI	
Jairam <i>et al.</i> ²³	480	Open AAA surgery or midline laparotomy in patients with BMI > 27 kg/m ²	Polypropylene	Onlay (188 patients)	24	Duration of surgery IH, postoperative complications	Clinical and ultrasound imaging or CT
						Quality of life Cost-effectiveness	
				Retromuscular (185 patients)			

IH, incisional hernia; SSI, surgical-site infection; DVT, deep vein thrombosis; AAA, abdominal aortic aneurysm; RYGB, Roux-en-Y gastric banding.



few years, several studies on the prevention of IH with prophylactic mesh reinforcement (PMR) have been conducted. Most RCTs included a limited number of patients. Different surgical techniques of PMR, including mesh placement in onlay, retromuscular or intraperitoneal position, have been studied. The European Hernia Society (EHS) Guidelines Development Group¹⁹ on the closure of abdominal wall incisions made a recommendation in

2015: that PMR to reduce the incidence of IH after elective midline laparotomy in a high-risk patient be suggested with a weak recommendation. The group also stated that larger trials were needed to make a strong recommendation.

Since the publication of these EHS guidelines, three meta-analyses^{20–22} have been published, together with the long-term data from the largest multicentre RCT on

Fig. 2 Forest plots of incisional hernia after prophylactic mesh reinforcement of a midline laparotomy versus primary suture**a Overall data****b According to risk of bias**

a Overall data from all 12 included studies; **b** data for studies with a low or high risk of bias. A Mantel-Haenszel random-effects model was used for meta-analysis. Risk ratios are shown with 95 per cent confidence intervals. IH, incisional hernia.

Fig. 4 Forest plots of incisional hernia after prophylactic mesh reinforcement of a midline laparotomy versus primary suture**a** Onlay mesh**b** Retromuscular mesh

a Data from studies using onlay mesh reinforcement; **b** data from studies using retromuscular mesh reinforcement. A Mantel–Haenszel random-effects model was used for meta-analysis. Risk ratios are shown with 95 per cent confidence intervals. IH, incisional hernia.

overall effect estimate did not overlap. The I^2 statistic was used to assess heterogeneity. Groups with zero events were adjusted with a constant continuity adjustment of 0.5 in each arm (as per the default adjustment in the software used). Publication bias was assessed by a funnel plot. Analyses were performed using Review Manager software (RevMan version 5.3; The Nordic Cochrane Centre, Copenhagen, Denmark). Two-sided $P < 0.050$ was considered statistically significant.


Conducting a meta-analysis can lead to type I errors (false-positives) or overestimation of treatment effects due to systematic errors (bias) and random errors (play of change). To avoid this, trial sequential analysis (TSA) was conducted. TSA can provide a required information size (RIS). The RIS is the required number of patients that needs to be included in the meta-analysis to provide firm evidence^{26,27}. Control event rate (CER) and relative risk reduction (RRR) values were calculated. CER is the proportion of participants in the control group who have the outcome. RRR can be interpreted as the reduction in the

relative risk of the specified outcome in the treatment group, compared with the control group. TSA was planned for all retrieved studies, and for the group of studies with low risk of bias. TSA was performed using the TSA software v0.9 (<http://www.ctu.dk/tsa/downloads.aspx>).

Results

A total of 1498 records were identified after removal of duplicates. After screening of title and abstract, 49 articles were found relevant for full-text assessment. After full-text assessment, 29 articles were excluded, leaving 13 RCTs and seven non-randomized trials for the qualitative synthesis (Fig. 1). In total, 13 RCTs^{5–11,23,28–32} fulfilled the inclusion criteria, but the study of Timmermans and colleagues³² was excluded because only the short-term results (postoperative complications in the first month) were discussed; thus 12 RCTs were analysed. Six studies^{7,8,11,23,29,31} were considered to have a low risk of bias, and six^{5,6,9,10,28,30} to have a high risk of bias (Table 1).



Fig. 5 Trial sequential analysis curves of the incidence of incisional hernia after prophylactic mesh reinforcement of a midline laparotomy versus primary suture



a Overall data from all 12 included studies; **b** data from studies with a low risk of bias. **a,b** Both graphs are two-sided. RIS, required information size.

The 12 RCTs comprised a total of 1815 patients. Study and patient characteristics are shown in *Table 2*. Inclusion criteria for PMR of the midline laparotomy in the individual RCTs were either the presence of an

AAA^{8,10,11,23}, morbid obesity^{5,7,9,30}, colorectal cancer surgery³¹ or a mixture of operative indications^{6,28,29}. Most studies placed a polypropylene mesh in an onlay^{6,10,23,29,31} or retromuscular^{7,8,11,23} position. Two studies^{10,30} used

Fig. 6 Forest plots of postoperative seroma after prophylactic mesh reinforcement of a midline laparotomy versus primary suture**a** Onlay mesh**b** Retromuscular mesh

a Data from studies using onlay mesh reinforcement; **b** data from studies using retromuscular mesh reinforcement. A Mantel–Haenszel random-effects model was used for meta-analysis. Risk ratios are shown with 95 per cent confidence intervals.

biological meshes, and one⁵ used a rapidly absorbable intraperitoneal mesh.



Outcome measurements

The primary outcome was the incidence of IH after follow-up of at least 12 months. Twelve RCTs were included in the overall quantitative analysis for the primary outcome and publication bias³³ was evaluated. The meta-analysis showed a significant reduction of IH in patients with PMR compared with that in patients who had a primary suture (RR 0.35; $P < 0.001$) (Fig. 2a). The funnel plot was slightly asymmetrical, indicating a possible publication bias for studies favouring mesh prophylaxis (Fig. 3). Analysis of the primary outcome for studies with a low risk of bias (6 RCTs) showed that the occurrence of IH was significantly lower (RR 0.23; $P < 0.001$) in the PMR group than in the primary suture group (Fig. 2b). Both onlay and

retromuscular PMR led to a significant reduction of the IH incidence compared with primary suture, with RRs of 0.26 ($P = 0.005$) and 0.28 ($P = 0.02$) respectively (Fig. 4).

TSA (for the primary outcome) was done for all 12 included studies. The CER proportion was 28 per cent, the RRR was 65 per cent, and a constant continuity adjustment was set at 0.5 events per group. The accrued information size ($n = 1815$) was 273.3 per cent of the estimated RIS ($n = 664$). This means that firm evidence was available. For low-risk-of-bias studies (6 RCTs), the CER proportion was 31 per cent and RRR 77 per cent. The accrued information size ($n = 1015$) was 283.5 per cent of the estimated RIS ($n = 358$) (Fig. 5).

Secondary outcomes were analysed for the low-risk-of-bias studies (6 RCTs). Patients with an onlay PMR had a higher risk of developing seroma (RR 2.23; $P = 0.030$) compared with patients who had a primary suture. Patients who had a retromuscular PMR had no

Fig. 7 Forest plots of surgical-site infection after prophylactic mesh reinforcement of a midline laparotomy versus primary suture**a** Onlay mesh**b** Retromuscular mesh

a Data from studies using onlay mesh reinforcement; **b** data from studies using retromuscular mesh reinforcement. A Mantel–Haenszel random-effects model was used for meta-analysis. Risk ratios are shown with 95 per cent confidence intervals. SSI, surgical-site infection.

greater chance of developing seroma than those with a primary suture (RR 1.67; $P=0.17$) (Fig. 6). The occurrence of SSI was not significantly higher for onlay PMR (RR 0.82; $P=0.33$) or retromuscular PMR (RR 0.85; $P=0.55$) than for primary suture (Fig. 7). There were insufficient data to analyse the incidence of haematoma and burst abdomen.

Discussion

This meta-analysis has shown that the use of PMR in patients undergoing midline laparotomy leads to a significantly lower occurrence of IH compared with a primary suture closure, with TSA indicating that the evidence was firm. This significant effect was shown for both onlay and retromuscular PMR. PMR was found to be safe, with no increase in SSI, but an increased risk of seroma formation for onlay PMR only.

This meta-analysis has a few limitations. Substantial statistical heterogeneity was seen in the studies regarding the primary outcome ($I^2=71$ per cent). This probably reflects

the variability of surgical technique and methodological approach across studies.

Although the technique of PMR in the treatment arm of the RCTs is often well described, there is less information on the control group undergoing primary suture. Protocols often describe the use of a suture:wound length ratio (SL:WL) greater than 4:1, but data for the SL:WL ratio were recorded and reported in only a few studies¹¹, so that adherence to the optimal primary suture technique was unclear in most studies. Moreover, the short-stitch technique, which is currently considered the best evidenced technique with the lowest incidence of IH³, was not used in any of the RCTs in this meta-analysis. Therefore, it could be argued that the treatment effect of PMR is increased because of a suboptimal suturing technique in the control groups.

Of the 12 RCTs included in this meta-analysis, half were considered to have a high risk of bias. Owing to the use of sensitivity analysis, however, the treatment effect was maintained, and was even greater when only studies with a low risk of bias were analysed.

Only RCTs with a minimum follow-up of 12 months were included in the meta-analysis, with all but one²⁹ having follow-up of at least 24 months. This is still too short, however, to evaluate the long-term efficacy or potential late adverse effects of PMR.

In most studies, physical examination was used to detect IH during follow-up. Some added selective or systematic medical imaging for the evaluation of IH, using either ultrasound imaging or CT. Imaging will increase the number of patients diagnosed with IH by detection of subclinical IHs, and this might overestimate the importance of PMR in providing clinical benefit.

Most of the 12 included studies used a polypropylene mesh in either an onlay or a retromuscular position; two^{10,30} used a biological mesh and one⁵ used an absorbable synthetic mesh. Two of these studies did not show PMR to be effective, and all three were considered to have a high risk of bias. All studies with a low risk of bias used either an onlay or a retromuscular mesh position, and all included trials were performed in an elective surgery setting. The results of this meta-analysis on PMR can therefore be considered valid only for synthetic, non-absorbable mesh in either an onlay or a retromuscular position in an elective setting.

The slightly asymmetrical funnel plot for IH indicates possible publication bias towards studies that favour PMR, resulting in an overestimation of the underlying beneficial effect of the intervention.

Most studies included only patients considered at high risk for developing an IH. It is difficult to identify the individual risk factors whereby patients would benefit from PMR. The guidelines on the closure of abdominal wall incisions from the EHS¹⁹ state that the evidence is weak for the use of PMR in patients at high risk of IH development. Such a guideline could be implemented only if the EHS Guidelines Development Group also described the exact criteria for considering patients to be at high risk. From this perspective, the study performed by Fischer *et al.*¹⁸ is interesting, as these authors stratified patients into four IH risk groups (low, moderate, high, extreme), based on characteristics of the patient and the surgical procedure. From this meta-analysis, it seems clear that patients undergoing AAA repair and those having bariatric surgery through a midline incision will benefit from PMR, although these are now less common as many patients with an AAA are treated with endovascular procedures, and bariatric surgery is often performed by a laparoscopic approach.

Selection based on patient characteristics needs cut-off values. The appropriate cut-off value for BMI is unclear. The PRIMA trial²³ used a BMI of at least 27 kg/m² as

an inclusion criterion for PMR, but further research is needed to identify an appropriate BMI cut-off point at which the risk of developing IH increases. Data from large clinically oriented prospective registries³⁴ might be helpful to explore which factors might lead to an increased risk of IH.

PMR can be considered safe for elective laparotomy. The only adverse event detected in this meta-analysis was an increased rate of seroma formation after onlay PMR, related to the subcutaneous dissection required. However, even though the occurrence of seroma was higher in the onlay PMR group, there was no increase in SSI. There were insufficient data to analyse the number of other adverse events, such as haematoma or burst abdomen.

One of the main strengths of this meta-analysis is the fact that RCTs with a low risk of bias were analysed separately³⁵. Further, the most up-to-date RCTs were included²³. TSA showed firm evidence in favour of PMR for midline laparotomy in high-risk patients, suggesting that no further trials are required to address the effects of PMR in this population. RCTs in other patient populations with differing levels of risk would, however, be helpful to evaluate the effectiveness of PMR.

Laparotomies are performed by a variety of surgical specialties, such as vascular surgeons, colorectal surgeons, gynaecologists and urologists. Many of those surgeons have little experience in treating abdominal wall hernias with mesh, particularly for retromuscular mesh placement. This meta-analysis has shown that onlay PMR, which is easier to perform, is also effective, and likely to be more acceptable to these surgeons.

This meta-analysis has provided evidence in favour of closure of midline laparotomies with PMR in high-risk patients. Individual risk factors should be taken into account to select patients who will benefit (most) from PMR, which should become standard treatment for high-risk groups.

Acknowledgements

A.P.J. and M.L.-C. contributed equally to this study.

Disclosure: The authors declare no conflict of interest.

References

- 1 Höer J, Lawong G, Klinge U, Schumpelick V. Factors influencing the development of incisional hernia. A retrospective study of 2983 laparotomy patients over a period of 10 years. *Chirurg* 2002; **73**: 474–480.
- 2 Mudge M, Hughes LE. Incisional hernia: a 10 year prospective study of incidence and attitudes. *Br J Surg* 1985; **72**: 70–71.

- 3 Millbourn D, Cengiz Y, Israelsson LA. Effect of stitch length on wound complications after closure of midline incisions: a randomized controlled trial. *Arch Surg* 2009; **144**: 1056–1059.
- 4 Bosanquet DC, Ansell J, Abdelrahman T, Cornish J, Harries R, Stimpson A *et al.* Systematic review and meta-regression of factors affecting midline incisional hernia rates: analysis of 14 618 patients. *PLoS One* 2015; **10**: e0138745.
- 5 Pans A, Elen P, Dewé W, Desaive C. Long-term results of polyglactin mesh for the prevention of incisional hernias in obese patients. *World J Surg* 1998; **22**: 479–482.
- 6 Gutiérrez de la Peña C, Medina Achirica C, Domínguez-Adame E, Medina Díez J. Primary closure of laparotomies with high risk of incisional hernia using prosthetic material: analysis of usefulness. *Hernia* 2003; **7**: 134–136.
- 7 Strzelczyk JM, Szymański D, Nowicki ME, Wilczyński W, Gaszynski T, Czupryniak L. Randomized clinical trial of postoperative hernia prophylaxis in open bariatric surgery. *Br J Surg* 2006; **93**: 1347–1350.
- 8 Bevis PM, Windhaber RA, Lear PA, Poskitt KR, Earnshaw JJ, Mitchell DC. Randomized clinical trial of mesh *versus* sutured wound closure after open abdominal aortic aneurysm surgery. *Br J Surg* 2010; **97**: 1497–1502.
- 9 Abo-Ryia MH, El-Khadrawy OH, Abd-Allah HS. Prophylactic preperitoneal mesh placement in open bariatric surgery: a guard against incisional hernia development. *Obes Surg* 2013; **23**: 1571–1574.
- 10 Bali C, Papakostas J, Georgiou G, Kouvelos G, Avgos S, Arnaoutoglou E *et al.* A comparative study of sutured *versus* bovine pericardium mesh abdominal closure after open abdominal aortic aneurysm repair. *Hernia* 2015; **19**: 267–271.
- 11 Muysoms FE, Detry O, Vierendeels T, Huyghe M, Miserez M, Ruppert M *et al.* Prevention of incisional hernias by prophylactic mesh-augmented reinforcement of midline laparotomies for abdominal aortic aneurysm treatment: a randomized controlled trial. *Ann Surg* 2016; **263**: 638–645.
- 12 Deerenberg EB, Harlaar JJ, Steyerberg EW, Lont HE, van Doorn HC, Heisterkamp J *et al.* Small bites *versus* large bites for closure of abdominal midline incisions (STITCH): a double-blind, multicentre, randomised controlled trial. *Lancet* 2015; **386**: 1254–1260.
- 13 van Ramshorst GH, Eker HH, Hop WC, Jeekel J, Lange JF. Impact of incisional hernia on health-related quality of life and body image: a prospective cohort study. *Am J Surg* 2012; **204**: 144–150.
- 14 van Ramshorst GH, Eker HH, van der Voet JA, Jeekel J, Lange JF. Long-term outcome study in patients with abdominal wound dehiscence: a comparative study on quality of life, body image, and incisional hernia. *J Gastrointest Surg* 2013; **17**: 1477–1484.
- 15 Gillion JF, Sanders D, Miserez M, Muysoms F. The economic burden of incisional ventral hernia repair: a multicentric cost analysis. *Hernia* 2016; **20**: 819–830.
- 16 Burger JW, Luijendijk RW, Hop WC, Halm JA, Verdaasdonk EG, Jeekel J. Long-term follow-up of a randomized controlled trial of suture *versus* mesh repair of incisional hernia. *Ann Surg* 2004; **240**: 578–583.
- 17 Helgstrand F, Rosenberg J, Kehlet H, Strandfelt P, Bisgaard T. Reoperation *versus* clinical recurrence rate after ventral hernia repair. *Ann Surg* 2012; **256**: 955–958.
- 18 Fischer JP, Basta MN, Mirzabeigi MN, Bauder AR, Fox JP, Drebin JA *et al.* A risk model and cost analysis of incisional hernia after elective, abdominal surgery based upon 12 373 cases: the case for targeted prophylactic intervention. *Ann Surg* 2016; **263**: 1010–1017.
- 19 Muysoms FE, Antoniou SA, Bury K, Campanelli G, Conze J, Cuccurullo D *et al.*; European Hernia Society. European Hernia Society guidelines on the closure of abdominal wall incisions. *Hernia* 2015; **19**: 1–24.
- 20 Borab ZM, Shakir S, Lanni MA, Tècce MG, MacDonald J, Hope WW *et al.* Does prophylactic mesh placement in elective, midline laparotomy reduce the incidence of incisional hernia? A systematic review and meta-analysis. *Surgery* 2017; **161**: 1149–1163.
- 21 Wang XC, Zhang D, Yang ZX, Gan JX, Yin LN. Mesh reinforcement for the prevention of incisional hernia formation: a systematic review and meta-analysis of randomized controlled trials. *J Surg Res* 2017; **209**: 17–29.
- 22 Payne R, Aldwinckle J, Ward S. Meta-analysis of randomised trials comparing the use of prophylactic mesh to standard midline closure in the reduction of incisional herniae. *Hernia* 2017; **21**: 843–853.
- 23 Jairam AP, Timmermans L, Eker HH, Pierik REGJM, van Klaveren D, Steyerberg EW *et al.*; PRIMA Trialist Group. Prevention of incisional hernia with prophylactic onlay and sublay mesh reinforcement *versus* primary suture only in midline laparotomies (PRIMA): 2-year follow-up of a multicentre, double-blind, randomised controlled trial. *Lancet* 2017; **390**: 567–576.
- 24 Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- 25 Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD *et al.*; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; **343**: d5928.
- 26 Thorlund K, Imberger G, Walsh M, Chu R, Gluud C, Wetterslev J *et al.* The number of patients and events required to limit the risk of overestimation of intervention effects in meta-analysis – a simulation study. *PLoS One* 2011; **6**: e2549.
- 27 Thorlund K, Engstrøm J, Wetterslev J, Brok J, Imberger G, Gluud C. *User Manual for Trial Sequential Analysis (TSA)*. Copenhagen Trial Unit, Centre for Clinical Intervention Research: Copenhagen, 2011.
- 28 El-Khadrawy OH, Moussa G, Mansour O, Hashish MS. Prophylactic prosthetic reinforcement of midline

- abdominal incisions in high-risk patients. *Hernia* 2009; **13**: 267–274.
- 29 Caro-Tarrago A, Olona Casas C, Jimenez Salido A, Duque Guileria E, Moreno Fernandez F, Vicente Guillen V. Prevention of incisional hernia in midline laparotomy with an onlay mesh: a randomized clinical trial. *World J Surg* 2014; **38**: 2223–2230.
 - 30 Sarr MG, Hutcher NE, Snyder S, Hodde J, Carmody B. A prospective, randomized, multicenter trial of Surgisis Gold, a biologic prosthetic, as a sublay reinforcement of the fascial closure after open bariatric surgery. *Surgery* 2014; **156**: 902–908.
 - 31 García-Ureña MÁ, López-Monclús J, Hernando LA, Montes DM, Valle de Lersundi AR, Pavón CC *et al.* Randomized controlled trial of the use of a large-pore polypropylene mesh to prevent incisional hernia in colorectal surgery. *Ann Surg* 2015; **261**: 876–881.
 - 32 Timmermans L, Eker HH, Steyerberg EW, Jairam A, de Jong D, Pierik EG *et al.* Short-term results of a randomized controlled trial comparing primary suture with primary glued mesh augmentation to prevent incisional hernia. *Ann Surg* 2015; **261**: 276–281.
 - 33 Schünemann H, Brozek J, Guyatt G, Oxman A (eds). *GRADE Handbook for Grading Quality of Evidence and Strength of Recommendations*; 2013. <http://gdt.guidelinedevelopment.org/app/handbook/handbook.html> [accessed 1 July 2017].
 - 34 Sedrakyan A, Campbell B, Graves S, Cronenwett JL. Surgical registries for advancing quality and device surveillance. *Lancet* 2016; **388**: 1358–1360.
 - 35 Roberts I, Ker K, Edwards P, Beecher D, Manno D, Sydenham E. The knowledge system underpinning healthcare is not fit for purpose and must change. *BMJ* 2015; **350**: h2463.

Supporting information

Additional supporting information can be found online in the Supporting Information section at the end of the article.